

[IMAGE]

Sleep Better, Naturally

By James P. Meschino, DC, MS

Chronic pain, which is reported to affect approximately 110 million Americans, is defined as three consecutive months of a painful condition. The most common conditions associated with chronic pain include arthritis / rheumatism; fibromyalgia; migraine headache; and low back pain. Evidence suggests that a multidisciplinary approach yields the best results in chronic pain management, whereas the method yielding the worst results for the patient, the health care system and society entails reliance on prescription narcotic drugs.

Over the years medical doctors have prescribed and recommended many analgesic drugs such as acetaminophen, nonsteroidal anti-inflammatory drugs (NSAIDs), and in more severe cases, narcotic drugs, as primary and sometimes exclusive methods of treatment in the management of chronic and acute muscle, joint and arthritic conditions. In recent years, documented evidence has shown that the frequent use of these medications for pain control has led to many serious unforeseen complications.

Fortunately, in recent years, research has shown the safe, effective pain-killing effects of California poppy (*Eschscholzia californica*). This herb has been shown to reduce night pain and induce sleep in patients with night pain without producing euphoria, addiction potential, physical dependency or serious side effects of any kind.

Health Complications From Standard Analgesic Drugs

Frequent use of acetaminophen has been shown to be a leading cause of liver failure, and acetaminophen ingestion is the leading cause of drug-induced liver failure, accounting for 50 percent of all acute liver failure cases in the U.S., half of which are unintentional (not suicide driven). Chronic intake of the recommended dosage of acetaminophen (up to 4 grams per day, with no single dose to exceed 1 gm) is responsible for most cases of acetaminophen-induced liver failure. Chronic use of acetaminophen has also been shown to damage the kidneys.

Heavy reliance on NSAIDs for chronic pain control has also yielded devastating health consequences. Recent studies confirm that in addition to gastrointestinal erosion, ulceration and bleeding, chronic NSAID

use also increases the risk of kidney damage, liver damage, congestive heart failure, high blood pressure and sudden cardiovascular death. Aspirin has long been associated with gastrointestinal damage and associated internal bleeding, but other NSAIDs are largely responsible for increased risk of cardiovascular death. This appears to be related to the promotion of thrombosis, associated with many NSAIDs from ibuprofen to diclofenac (Voltaren) to COX-2 inhibitors (e.g., Celebrex, Vioxx)

As such, doctors have been instructed not to recommend any NSAIDs, other than aspirin, for patients at high risk for heart disease. These recommendations also extend to precluding the recommendation of all NSAIDs for patients with any compromised kidney function. Low-dose aspirin, although recommended as a blood thinner for those who have suffered a previous heart attack, is no longer recommended to prevent first heart attack (primary prevention) due to the increasing reports of intestinal bleeds and bleeding into the brain, seen in patients prescribed low-dose aspirin (75-81 mg) for this purpose.

Narcotic Drugs – Rising Concerns About Addiction

Since the early 1990s governments have allowed doctors to prescribe narcotic drugs (e.g., oxycodone) for patients presenting with a wide variety of musculoskeletal pain conditions. Prior to this, narcotic drugs were only prescribed for patients with intractable pain, primarily due to terminal cancers (e.g., morphine drip). As such, physicians commonly use narcotics to reduce a patient's post-operative pain or to reduce anxiety and induce anesthesia prior to an operation. These drugs are also commonly prescribed in an attempt to enable individuals with chronic pain to lead productive lives.

The problem is that many people who are prescribed and taking opioids for a period of time develop a physical dependence on the drug which can lead to abuse of the painkiller. Studies now show that 2.5 million Americans, of the 4.7 million who begin to abuse prescription drugs in any given year, use pain pills. Thus, more than 50 percent of all drug abuse cases involve analgesic drugs, and very often narcotics.

Recognizing the potential for opioid abuse, addiction, diversion and related mortality, many jurisdictions have developed guidelines or implemented programs to promote more judicious use of these drugs. Across the board, medical doctors are being instructed to cut back on their prescription writing for narcotic drugs, and systems are being put in place to track and integrate pharmacy dispensing of these drugs using electronic recording and monitoring systems.

A Safe Herbal Alternative

Recent studies have shown that the medicinal ingredients in the herb *Eschscholzia californica* (California poppy) block nighttime pain, allowing the patient to sleep through the night without being awakened by musculoskeletal pain. The herb also helps to induce sleep, enabling patients who are in pain to fall asleep and experience a restful sleep through the night. This, in turn, allows more rapid healing and improved response to other treatments.

The active ingredients in *Eschscholzia californica* relieve pain without producing euphoria or having addiction potential. Stimulation of opioid receptors blocks pain sensation in the brain and blocks pain conduction in the spinal cord from reaching higher brain centers. Activation of serotonin receptors is also known to block the sensation of pain and induce sleep.

Unlike narcotic drugs (e.g., Percocet, Oxydone) and benzodiazepine drugs (e.g., Valium, Ativan) often used to help patients in pain sleep through the night, supplements containing *Eschscholzia californica* do not cause addiction or destroy a person's motivation to return to a productive life. The active constituents in this herb do not cause euphoria or feeling of being "stoned," which allows individuals to function normally and better comply with treatment recommendations, including exercise.

Precautionary Notes

Patients should not take this herb if they are taking an evening or nighttime dose of a narcotic drug (e.g., Percodan, Oxycontin), anti-anxiety drug and/or a sleep-inducing drug (e.g., Valium, Sonata, Ambien). Patients taking narcotic or benzodiazepine drugs who wish to wean themselves off of these drugs by using *Eschscholzia californica* as a replacement for chronic pain management, must do so under the supervision and monitoring of their attending physician. Narcotic and benzodiazepine drugs are highly addictive; thus, each case requires individualized evaluation and attention. As always, talk to your doctor for more information.

James Meschino, DC, MS, practices in Toronto, Ontario, Canada and is the author of four nutrition books, including *The Meschino Optimal Living Program* and *Break the Weight Loss Barrier*.

Page printed from:

http://www.toyourhealth.com/mpacms/tyh/article.php?id=1741&no_paginate=true&no_b=true