

[IMAGE]

## Stand Up Straight

By Dr. Jeffrey Tucker

I remember reading a quote in the *American Journal of Pain Management* that said "To live a long, active, energetic life, few things matter more than good posture." Postural issues are a big contributor to many different aches and pains and injuries to our bodies. Injuries related to poor posture tend to be overuse injuries, which build up over a period of time. Slouched sitting for extended periods of time at a desk or in front of the TV can cause the shoulder joints to sit in a forwards position. This causes a muscle imbalance where the chest muscles are tight and the upper back muscles are weak. If you suffer with low back pain that developed slow and gradual with no history of trauma or overuse, the problem may be due to poor posture. Slumped sitting usually causes the arch of the back to flex or round and this places extra strain on the muscles and ligaments, which support the lower back. This results in muscle spasms and sometimes muscle strains.

Sitting, staring at a computer screen for hours on end, allowing your shoulders to round and your neck protrudes forward can cause aches and stiffness in the neck-shoulder area and even cause headaches. An accumulation of poor posture day-in and day-out can result in shortening of the chest muscles and weakening of the small, postural upper back and neck muscles, which work to pull the shoulders back. Once the rounded shoulders and forward head posture become a habit, it is hard to break that pattern.

lower back pain - Copyright © Stock Photo / Register Mark Most people get out of bed in the morning and go sit down at a table and eat breakfast, then get in their car and drive to work. Large chunks of the work day is spent sitting hunched over a computer or in a vehicle driving to appointments. After work people go home and sit at the dinner table and then sit slumped on a couch to watch TV until they go to bed. This excess sitting for long periods during the day and night adversely affects posture, which in turn affects your performance in your activities and is quite often a predisposing factor in injury.

I understand most people need to spend on average eight to 10 hours each day at work. Don't be one of those people who sit unconsciously in improper body positions and engage in repetitive movements that create muscle imbalances leading to poor posture. Poor self esteem, psychological distress & depressive symptoms are all related to poor posture. The most natural thing you can do here is increase your "get up"

and "move around" time. Create a variety of movement in your activities of daily living.

If you have poor flexibility, try some simple yoga. Muscle imbalances and joint dysfunctions associated with poor posture can create areas of too much motion in certain spinal segments causing instability. These areas may then wear out prematurely, while other areas may have too little motion in the spine causing range of motion/mobility dysfunctions; anytime you have a right side - left side imbalance, we call that an asymmetry. If you have an asymmetry in your muscles, you are more susceptible to injury.

Improve your posture by using these techniques:

1. Become aware of the things that you are doing, even the things that you don't even know you are doing that are contributing (harming) to your posture.
2. Think of staying in a "tall spine" posture (while sitting, standing, during exercise).
3. Take frequent breaks from sitting and use the Brugger's postural relief position as one of your style of breaks.
4. Know what it feels like to be in proper posture alignment and frequently try to duplicate that feeling - sometimes clients don't even know what good posture feels like and looks like.

Taking frequent breaks from sitting at your desk is one of the most important things you can do for prevention of poor posture. Become aware of the times that you are doing repetitive movements and/or sustained postures, i.e., the mattress you sleep on may be worn out and contribute to microtrauma to the tissues causing altered spinal curves. The position you sleep in is important - the least offensive sleep position is on your back, then side lying with a pillow between the knees, and the least desirable position is on the stomach. A pillow with a good cervical support is important - a pillow without any cervical support may contribute to altered neck curves. The chair at your work station should allow you to sit upright rather than in a slumped posture.

Other things that maybe harming our posture: I think our moods influence our posture; a person who is depressed has a classic hunched over looking appearance. Even our exercise choices need to be scrutinized. If you perform the same exercise over and over such as cyclists who spend 2-3 hours riding their bicycles in a position of lumbar flexion develop a reduced lumbar curve; long distance swimmers who perform repeated motions may experience shoulder pain from altered posture and faulty biomechanics. For any person who sits eight hours a day hunched over a computer, the last thing that person needs to do is spend time hunched over a bicycle for recreation or pounding out bench presses at the gym.

The shoes you wear daily are important to maintain - worn out soles could contribute to foot and ankle malpositions leading to altered posture; foot pronation issues may require an insert or orthotic - this can help improve gait and posture by correcting faulty biomechanics.

I always recommend that we improve our ability to take deep breaths and expand the lungs capacity. Using the cue "breath into the back" helps improve posture.

Let me be perfectly clear – you can improve your posture – first become aware of your posture. Second requires training your body with simple exercise maneuvers and progressing to more challenging strength exercises.

Here are some simple exercises to get you started:

1. Engage in daily use of the foam roll to provide self-myofascial release and self massage. Spend 3-5 minutes rolling out the thoracic spine and shoulders.
2. Make sure you know how to go from "sitting to standing" properly. Stand upright (tall spine) imagining a string attached to the base of the skull is lifting you upright, rather than leaning forward at the waist when going from sitting to standing. Once you are up, raise the hands above the head with the arms extended and with the elbows in line with the ears. Lean or bend backward as far as possible, making sure the hips go forward and the arms go backwards simultaneously. Repeat this maneuver 10 times.
3. Perform "Chair Decompression": The person sits in an upright chair with their arms behind them, slightly bent, hands on the seat of the back of the chair. They push downward, straightening the arms and leaving the buttocks in the chair, unloading the trunk and spine. Keep the arms externally rotated; this moves the upper body into something similar to Brugger's.
4. Perform Brugger's relief position: Sit at the edge of a chair; Put your knees apart (wide) and your feet under the knees: Arch your back; Rotate your arms outward so your palms face forward; Separate your fingers and point your thumb backward; Tuck in your chin; Hold this position while taking a deep breath in through your abdomen. HOLD the position for 5 seconds, release for 3 seconds, Repeat 3-5 times.
5. Perform Cobra: Laying face down on the floor-in prone position, have arms beside your hips. Activate the core by drawing in your navel towards spine and squeezing the glutes. With your core and glutes activated, lift the chest off the floor, lift arms up and back towards the hips rotating thumbs towards the ceiling. Pause momentarily at the top of the lift then return to starting position; at all times keeping the

chin tucked into the chest and the feet on the floor. Upon completion of the movement, repeat. Don't over emphasize arching of the back to the lift the chest off floor. Only lift to where you are comfortable - no lower back pain should be felt. Note: hold for 2-3 seconds. Repeat 5 times.

6. Core training including the abdominals, lower back, gluteus, and hips is important for pelvis alignment.
  7. Strength training exercises include A) Bent over back rows. Bend over from the hips with the torso parallel to the floor. Pull either bands or free weights up, squeezing your shoulder blades as close together as you go). B) Standing or seated rowing exercises - start with your arms in front of the body holding on to a band or cable machine. Pull straight back bending at the elbows with the hands moving back along the sides of the body. C) Back Flys - Gripping on to a cable machine or bands, extend your arms into a wing span position.
- 

**Jeffrey Tucker, DC**, is a doctor of chiropractic and rehabilitation specialist who integrates chiropractic, exercise and nutrition into his practice in West Los Angeles. You can sign up for his newsletter at [DrJeffreyTucker.com](http://DrJeffreyTucker.com).

Page printed from:

[http://www.toyourhealth.com/mpacms/tyh/article.php?id=1497&pagenumber=1&no\\_paginate=true&no\\_b=true](http://www.toyourhealth.com/mpacms/tyh/article.php?id=1497&pagenumber=1&no_paginate=true&no_b=true)